

Diffusion in a generalized Rubinstein-Duke model of electrophoresis with kinematic disorder.

Richard D. Willmann* and Gunter M. Schütz

Institut für Festkörperforschung, Forschungszentrum Jülich, 52425 Jülich, Germany

Kavita Jain

Department of Theoretical Physics, Tata Institute of Fundamental Research,

Homi Bhabha Road, Mumbai 400 005, India

(Dated: 2nd February 2008)

Abstract

Using a generalized Rubinstein-Duke model we prove rigorously that kinematic disorder leaves the prediction of standard reptation theory for the scaling of the diffusion constant in the limit for long polymer chains $D \propto L^{-2}$ unaffected. Based on an analytical calculation as well as Monte Carlo simulations we predict kinematic disorder to affect the center of mass diffusion constant of an entangled polymer in the limit for long chains by the same factor as single particle diffusion in a random barrier model.

PACS numbers: 61.25.Hq, 83.10.Kn

The derivation of large scale properties of polymer systems such as viscosity and diffusion constant out of microscopic models is among the basic problems of polymer science. For entangled polymers in gels or melts, de Gennes¹ predicted scaling laws for the dependence on the polymer length L for viscosity ($\eta \propto L^3$) and diffusion constant ($D \propto L^{-2}$). These scaling laws are assumed to be valid for the limiting case of polymer length going to infinity. In experiments, the apparent scaling laws $D \propto L^{-2.4}$ and $\eta \propto L^{3.4}$ are found^{2,3,4}. However, the experimental findings do not contradict the predictions of reptation, as a crossover due to decreasing finite size effects with increasing chain length cannot be ruled out³. Presumably, contour length fluctuations (CLF) are one cause for the deviating scaling exponent for finite chain length^{5,6,7}. In the framework of the repton model introduced by Rubinstein⁸ and Duke^{10,11} (further on to be called RD model), which incorporates CLF, it is possible to calculate viscosity, diffusion coefficient and other quantities of interest. Good agreement both with theoretical and experimental results is found^{6,7,14,15}. Using the RD model with periodic boundary conditions Kooiman and van Leeuwen¹² analytically calculated the proportionality constant c for the diffusion constant in the limit for infinite chain length: $\lim_{L \rightarrow \infty} DL^2 = c$. They found $c = w/(2d + 1)$, d being the dimensionality of the entanglement network and w a model constant defining the time scale. Building upon this result, Prähofer and Spohn¹³ rigorously derived the leading order term in $1/L$ for the diffusion constant in the RD model and furthermore proposed a scaling for the finite size effects: $DL^2 - W/(2d + 1) \propto L^{-\beta}$, where $1/2 \leq \beta \leq 1$. All results mentioned above were obtained under the assumption that the entanglement network which is topologically restricting the polymer is regular and static. Real entanglement networks such as gels or melts are disordered. The effects of a disordered environment can be manifold:

- Spatial variations of the mobility of the 'defects' of stored length.
- Locally fluctuating potential energy due to interactions between chain and environment.
- Entropically favorable regions of low entanglement density.
- Relaxation of the environment (constraint release).

Numerical investigations¹⁶ showed that entropically favorable regions can for short chains substantially lower the diffusion constant by the creation of 'entropic traps'. So far, conclu-

sive investigations of diffusing polymers long enough to span several such traps are missing. Constraint release is considered to be of minor importance in gels but needs to be self consistently taken into account in melts¹⁷. Schäfer, Baumgärtner and Ebert¹⁹ numerically investigated the effect of kinematic disorder, i.e., disorder which affects the mobility of the chain segments while leaving the equilibrium distribution of chain configurations unaltered. Their investigation shows that reptation prevails in presence of kinematic disorder. However, due to being based on MC simulations and thus relatively short chains no quantitative results could be obtained about the modifications the disorder would cause to the reptation prediction for the diffusion constant in the long chain limit. As reptation is shown to prevail, the scaling $\lim_{L \rightarrow \infty} DL^2 = c$ must remain valid. The aim of this article is to calculate the coefficient c for a polymer diffusing in a disordered environment exhibiting kinematic disorder. This means that we are interested in the behavior of the chain in the long chain length limit originally envisaged by standard reptation theory. Some of the results presented here in detail were briefly reported in an earlier communication¹⁸.

The article is divided into six chapters: In chapter I the definition of the model is given and the master equation for the chain dynamics is presented in terms of the quantum Hamiltonian formalism²⁰. This master equation yields the stationary distribution of chain conformations as shown in chapter II. The model with periodic boundary conditions is analyzed in chapter III. Adapting the strategy in Ref. 12 for obtaining the diffusion constant for the periodic system yields a lower bound on the diffusion constant for the original system. Chapter IV is devoted to deriving an upper bound to the diffusion constant by generalizing a variational approach used in Ref. 13 for the ordered system. In chapter V Monte Carlo simulations of the polymer diffusion in kinematically disordered environments are presented. The last chapter addresses the dynamics of internal chain segments.

I. A LATTICE MODEL FOR REPTATION WITH KINEMATIC DISORDER

The RD model is a discretized model for reptation, i.e., it describes the dynamics of a polymer in an entanglement network. This network is assumed to be regular and static. In 3 dimensions it has the shape of a cubic lattice, where the polymer is forbidden to cut through the edges of the cubes. The faces of the cubes can be penetrated by the chain (see figure 1). The polymer itself is assumed to consist of $L + 1$ 'reptons', i.e., segments of about

the entanglement length, which equals the lattice constant of the cubic lattice.




Figure 1: Two dimensional representation of a kinematically disordered lattice with a reptating polymer chain and the lattice gas mapping. Arrows indicate possible moves of the reptons. $B = \exp(E/2)$.

The model defines dynamical rules for the reptons' motion. The dynamics on a smaller scale, i.e. high frequency Rouse modes, is averaged out. In this sense, the RD model is an effective model. The dynamical rules for the reptons are:

- Each cell occupied by the chain must contain at least one repton to ensure connectivity of the chain, since the length represented by a repton equals the lattice constant of the cells. The sequence of occupied pores corresponds to the tube as in standard reptation theory.
- End reptons can move to adjacent cells provided rule a) is not violated.
- Interior reptons can move to cells occupied by the neighboring reptons if allowed by rule a). This ensures the dynamics to be reptation, as any movements in the interior requires multiple reptons to occupy the same cell, which means that there is some excess 'stored length' available. This corresponds to the motion of 'defects' in the original de Gennes reptation model.

All moves are assumed to be thermally activated. In our model the cubic lattice representing the entanglement network gets kinematically disordered by assigning an individual energy barrier to every boundary between adjacent cells. These barriers have to be overcome by a crossing repton. Note that we do not assign different energies to the reptons when residing in the cells. This choice of quenched disorder guarantees that only the mobility of the reptons, but not the equilibrium configurations are affected, as will be seen below. Let there be $\sigma \in N$ different possible hopping rates W_α , which are distributed randomly and occur with a probability $f(W_\alpha)$ throughout the lattice. We demand that the disorder averages $\langle 1/W \rangle = \sum_{\alpha=1}^{\sigma} f(W_\alpha)/W_\alpha$ and $\langle 1/W^2 \rangle$ (the mean hopping time and its second moment) are finite.

The RD model is used to describe the dynamics of an entangled polymer chain under the influence of an external electric field. A common example is DNA under electrophoresis conditions where the reptons carry a charge each and develop a drift velocity along the direction of an applied electric field. Let the field be oriented along the (111) diagonal of the cubes. We denote the (dimensionless) energy gain of a repton when moving from one cell to another along the direction of the field as E . By local detailed balance, moves across a cell boundary with assigned hopping rate W_α along the field happen with rate $W_\alpha \exp(E/2)$, those in the opposite direction with rate $W_\alpha \exp(-E/2)^{10}$. By projecting the reptons' positions on the direction of the field, their relative coordinates can be denoted as a 1-d lattice gas with L sites by the following prescription:

- If the projected link between adjacent reptons i and $i + 1$ is oriented along (against) the field direction across a cell boundary with assigned rate W_α , site i of the lattice gas is assigned the value $y_i = \alpha$ ($y_i = -\alpha$). We interpret values $\pm\alpha$ as particles.
- If adjacent reptons i and $i + 1$ occupy the same cell, i.e., their projected positions coincide, the link is represented in the lattice gas as $y_i = 0$ at site i . We interpret a site, which is assigned a 0 as being unoccupied.

Thus the $L + 1$ coordinates of the reptons in direction of the field are translated into the equivalent set of the relative coordinates manifest as the assignment of $\alpha, -\alpha$ or 0 to the L sites of the lattice gas plus the center of mass coordinate's component in field direction. The dynamics of the reptons translates into the lattice gas picture as follows: In

the bulk particles of sort ' $\pm\alpha$ ' hop to the left with rate $W_\alpha \exp(\pm E/2)$ and to the right with rate $W_\alpha \exp(\mp E/2)$, where each site can be occupied by at most one particle. The end dynamics in the lattice gas picture needs some care: Assuming y_1 (y_L) to be non zero, the only possible move is the retraction of the end repton to the cell occupied by it's neighbor (rule a)). This retraction, being a particle annihilation event in the lattice gas picture, happens with the same rate as the respective move in the bulk. Assuming y_1 (y_L) to be zero, the end repton can, according to rule b), move to any of the $2d$ adjacent cells. For half of these the move leads to links being along the field direction, the other half against it. The probability of the chosen move leading to the crossing of a cell boundary with rate W_α being assigned to it is $f(W_\alpha)$. Thus the move of the repton, being a particle creation event in the lattice gas picture leads to y_1 (y_L) changing from '0' to ' $\pm\alpha$ ' with rate $\exp(\mp E/2)f(W_\alpha)W_\alpha d$ ($\exp(\pm E/2)f(W_\alpha)W_\alpha d$). This choice of boundary dynamics is on average correct, but neglects the actual local structure of the network. In order to verify that the model is correct on long time scales (as relevant for the CMS diffusion) we performed Monte Carlo simulations comparing the diffusion constant of our projected model with $d = 2$ at different disorder distributions with a polymer chain moving according to the dynamics of the repton model in a two dimensional lattice with random but fixed hopping rates assigned to each cell boundary. It turns out that although the results differ for small chains ($L < 10$) they coincide within the statistical errors for longer chains. This legitimates our choice of boundary dynamics for investigating the behavior of long chains. Figure 2 shows an example where a disorder distribution with $\langle W \rangle = 0.505$ was chosen.

Every move of a particle in the lattice gas leads to a change in the component of the center of mass coordinate of the repton chain along the field direction.

- Particle type ' α ' moving to the right (left) decreases (increases) x by $1/(L+1)$, as this is equivalent to a repton moving downward (upward). As there are $L+1$ reptons each contributes $1/(L+1)$ to the center of mass position.
- Particle type ' $-\alpha$ ' moving to the right (left) increases (decreases) x by $1/(L+1)$.

In the subsequent chapters we will calculate the drift velocity v of the center of mass coordinate in presence of an applied electric field and by employing the Einstein relation the




Figure 2: Comparison between the diffusion coefficients for simulations of the disordered RD and two dimensional lattice model.

zero field diffusion constant. When calculating v we restrict ourselves to the linear response regime, ignoring higher order field dependences. In order to calculate v only the change in the center of mass coordinate along the field x has to be known, but not the absolute value itself. Whereas the latter cannot be known from the lattice gas, the former is given by the difference of the currents of positive particles j^+ and that of the negative ones j^- : $v = j^- - j^+$. As the choice of the field direction is arbitrary, by use of the Einstein relation the model allows for computing the zero field diffusion constant along a distinct direction in d -dimensional space. As zero field diffusion is isotropic this immediately yields the d -dimensional diffusion constant. This is due to distinguishing between particles of types α and $-\alpha$ in contrast to the original projected repton model used by Rubinstein, which allows only for the computation of the curvilinear diffusion constant along the contour of the chain. In the Rubinstein model additional assumptions are necessary to relate the curvilinear to the d -dimensional diffusion constant. Our model as well as the RD model allows for computation of the latter quantity within the model.

II. QUANTUM HAMILTONIAN AND STATIONARY STATE

The model introduced in the previous chapter describes a Markov process and thus the dynamics can be written in form of a master equation. For convenience we will use the

quantum Hamiltonian formalism²⁰ to write down the master equation at zero electric field and solve for the stationary state.

At each site of the lattice gas with length L , $2\sigma + 1$ values for y_i are possible. Therefore, the state space X has the dimension $(2\sigma + 1)^L$. Every $\eta \in X$ is assigned a vector $|\eta\rangle$ and a transposed vector $\langle\eta|$. These vectors constitute a basis for the space of system configurations X . A probability distribution $P(\eta) \equiv P_\eta$ can thus be written as a probability vector:

$$|P\rangle = \sum_{\eta \in X} P(\eta) |\eta\rangle. \quad (1)$$

Let a summation vector $\langle s|$ be defined as $\langle s| = \sum_{\eta \in X} \langle\eta|$. Normalization of the probability vector is given if $\langle s|P\rangle = 1$. The generator for the dynamics of the system is the matrix H : The off-diagonal elements of the matrix contain the (negative) transition rates $t(\eta, \eta')$ from a state η' to η :

$$\langle\eta|H|\eta'\rangle = H_{\eta,\eta'} = -t(\eta, \eta'). \quad (2)$$

The diagonal elements $H_{\eta,\eta}$ contain the sum of all outgoing rates :

$$\langle\eta|H|\eta\rangle = H_{\eta,\eta} = \sum_{\eta' \neq \eta} t(\eta', \eta). \quad (3)$$

It is easily checked that conservation of probability, i.e. $\langle s|H = 0$ in the language of the quantum Hamiltonian formalism, is fulfilled by H . The master equation

$$\frac{d}{dt}P_\eta(t) = \sum_{\eta' \neq \eta, \eta \in X} \left(t(\eta, \eta')P_{\eta'}(t) - t(\eta', \eta)P_\eta(t) \right) \quad (4)$$

describing the Markovian dynamics of the system can be written as:

$$\frac{d}{dt}|P(t)\rangle = -H|P(t)\rangle. \quad (5)$$

The stationary state $|P^*\rangle$ is thus characterized by the equation

$$H|P^*\rangle = 0. \quad (6)$$

In this formalism the expectation value $\langle F(t) \rangle = \sum_{\eta} F(\eta) P(\eta, t)$ of an operator F is written as follows²⁰: The operator $F : X \rightarrow X$ is represented by a diagonal matrix $F = \sum_{\eta} F(\eta) |\eta\rangle\langle\eta|$. Then

$$\begin{aligned} \langle F(t) \rangle &= \langle s | F e^{-Ht} | P(0) \rangle \\ &= \langle s | e^{Ht} F e^{-Ht} | P(0) \rangle. \end{aligned} \quad (7)$$

Let the time-dependent operator $F(t)$ for $t > 0$ be defined as

$$F(t) = e^{Ht} F e^{-Ht}. \quad (8)$$

$\langle F(t) \rangle$ is an expectation value that it is not only averaged over possible realizations of the process but also over the initial states according to $P(0)$. In the following it is assumed that $|P(0)\rangle = |P^*\rangle$, so that

$$\langle F(t) \rangle = \langle s | F(t) | P^* \rangle. \quad (9)$$

For our model, we choose a tensor product basis as follows: Let at each site of the lattice gas the unit vector e_1 denote $y_i = 0$, $e_{2\alpha}$ denote $y_i = \alpha$ and $e_{2\alpha+1}$ denote $y_i = -\alpha$. A state vector for a state $\eta = (1, -3, \dots, 5, -2)$ for example can then be written as $|\eta\rangle = e_2 \otimes e_7 \otimes \dots \otimes e_{10} \otimes e_5$. Using this basis, the following operator creates a particle of type α at site i , provided it was previously unoccupied:

$$a_{\alpha,1}^\dagger(i) = \underbrace{\mathbf{1} \otimes \dots \otimes}_{i-1} E_{(2\alpha,1)} \underbrace{\otimes \dots \otimes \mathbf{1}}_{L-i} = E_{(2\alpha,1)}(i), \quad (10)$$

where $E_{(2\alpha,1)}$ is the matrix with a single entry 1 at row 2α and column 1. Similarly, the operator $a_{\alpha,-1}^\dagger(i) = E_{(2\alpha+1,1)}(i)$ creates a particle of type $-\alpha$, if possible. The corresponding annihilation operators at site i are $a_{\alpha,1}(i) = E_{(1,2\alpha)}(i)$ and $a_{\alpha,-1}(i) = E_{(1,2\alpha+1)}(i)$. To formulate the diagonal part of the quantum Hamiltonian the matrices $u(i) = E_{(1,1)}(i)$, $v_{\alpha,1}(i) = E_{(2\alpha,2\alpha)}(i)$ and $v_{\alpha,-1}(i) = E_{(2\alpha+1,2\alpha+1)}(i)$ are employed. Thus, the quantum Hamiltonian for the model defined in the previous chapter at zero field reads:

$$\begin{aligned}
H_{open} = & \sum_{i=1}^{L-1} \sum_{s=\pm 1} \sum_{\alpha=1}^{\sigma} W_{\alpha} [-a_{\alpha,s}^{\dagger}(i+1)a_{\alpha,s}(i) - a_{\alpha,s}^{\dagger}(i)a_{\alpha,s}(i+1) \\
& + u(i+1)v_{\alpha,s}(i) + u(i)v_{\alpha,s}(i+1)] \\
& + \sum_{s=\pm 1} \sum_{\alpha=1}^{\sigma} [W_{\alpha} df(W_{\alpha})(-a_{\alpha,s}^{\dagger}(1) + u(1)) + W_{\alpha}(-a_{\alpha,s}(1) + v_{\alpha,s}(1))] \\
& + \sum_{s=\pm 1} \sum_{\alpha=1}^{\sigma} [W_{\alpha} df(W_{\alpha})(-a_{\alpha,s}^{\dagger}(L) + u(L)) \\
& + W_{\alpha}(-a_{\alpha,s}(L) + v_{\alpha,s}(L))]
\end{aligned} \tag{11}$$

Plugging a product measure ansatz

$$|P_{open}^*(0)\rangle = \begin{pmatrix} 1 \\ p_{1,1}(1) \\ p_{1,-1}(1) \\ p_{2,1}(1) \\ \vdots \end{pmatrix} \otimes \dots \otimes \begin{pmatrix} 1 \\ p_{1,1}(L) \\ p_{1,-1}(L) \\ p_{2,1}(L) \\ \vdots \end{pmatrix} \frac{1}{\prod_{i=1}^L (1 + \sum_{\alpha=1}^{\sigma} (p_{\alpha,1}(i) + p_{\alpha,-1}(i)))} \tag{12}$$

into $H_{open}|P_{open}^*(0)\rangle = 0$ leads to a very simple set of equations for the probabilities $p_{\alpha,\pm 1}$ and one finds

$$|P_{open}^*(0)\rangle = \left(\begin{pmatrix} 1 \\ df(W_1) \\ df(W_1) \\ \vdots \\ df(W_{\sigma}) \\ df(W_{\sigma}) \end{pmatrix} \right)^{\otimes L} \frac{1}{(2d+1)^L}. \tag{13}$$

The geometrical equilibrium conformation of the chain depends on the probability of occurrence for links between reptons along or against the field, irrespective of the assigned hopping rates of possibly crossed cell boundaries. This means we have to consider the overall probability for particles of positive sign at a site, which is $\sum_{\alpha=1}^{\sigma} f(W_{\alpha})d/(2d+1) = d/(2d+1)$, or negative sign, being also $d/(2d+1)$, respectively. These are equal to the probabilities found for the disorder free RD-model²² which shows that the chosen kind of disorder is indeed

kinematic disorder, as it influences only the mobility but not the equilibrium configurations of the chain.

Note that it is not possible to compute the stationary state for the model with a non-zero field at arbitrary L . Therefore the Einstein relation $1/L \, d/dE(< v(E) >) = D$ is not a feasible way of straightforwardly computing the diffusion constant. Extra input is needed to overcome this problem.

III. MODEL WITH PERIODIC BOUNDARY CONDITIONS

In this section we adapt a calculation due to Kooiman and van Leeuwen¹² for the original RD-model with periodic boundary conditions to the case with kinematic disorder. The quantum Hamiltonian for the periodic system at zero field reads

$$H_{per} = \sum_{s=\pm 1} \sum_{\alpha=1}^{\sigma} \sum_{i=1}^L W_{\alpha} \left[-a_{\alpha,s}^{\dagger}(i+1)a_{\alpha,s}(i) - a_{\alpha,s}^{\dagger}(i)a_{\alpha,s}(i+1) + u(i+1)v_{\alpha,s}(i) + u(i)v_{\alpha,s}(i+1) \right]. \quad (14)$$

Plugging $|P_{open}^*(0) \rangle$ as for the open system into the equation $H_{per}|P_{open}^*(0) \rangle = 0$ for the periodic system shows that at zero electric field the stationary state of the open system is also stationary with respect to the dynamics of a corresponding periodic system. In a periodic system, the phase space is non ergodic, as neither the order nor the number of occurring particles on such a ring can be changed. Therefore, every connected subset of the phase space ('channel') has its own stationary state. We can calculate the stationary state for the periodic system in presence of a field E by mapping the system to a disordered zero range process, as introduced by Benjamini, Ferrari and Landim²³. This means that the Einstein relation can be employed to obtain the diffusion constant in the periodic case. Here, the definition of the drift velocity of the center of mass and the corresponding diffusion constant are induced from the system with open boundary conditions: $v = j^- - j^+$.

Instead of characterizing the system by the spins on the lattice gas $\mathbf{y} = (y_1, \dots, y_L)$ it can equivalently be characterized by the sets $\mathbf{s} = (s_1, \dots, s_L)$ containing the signs of the non zero y_i , $\mathbf{w} = (w_1, \dots, w_L)$ containing the absolute values of these y_i and $\mathbf{n} = (n_1, \dots, n_L)$ where n_i amounts to the number of '0' between s_i and s_{i+1} on the lattice gas. Thus every lattice gas configuration on a ring can be translated into a configuration on a lattice of length M

in the zero range picture: Site i carries n_i zero range particles and is separated from site $i + 1$ by a bond characterized by s_{i+1} and w_{i+1} . The total number K of zero range particles must equal the total number $L - M$ of '0' in the lattice gas: $\sum_{i=1}^M n_i = K = L - M$. The dynamics of the lattice gas picture translates into the zero range picture as follows: The configuration $(\dots, n_j, n_{j+1}, \dots)$ changes to $(\dots, n_j - 1, n_{j+1} + 1, \dots)$ with rate $h_{j+1}^{-1} w_{j+1}$ and to $(\dots, n_j + 1, n_{j+1} - 1, \dots)$ with rate $h_{j+1} w_{j+1}$, where $h_j = \exp(-Es_j/2)$. This means that the random hopping rates as well as the s_j are not assigned to individual particles, but to bonds between sites in the zero range (zr) picture. At $E = 0$ the zr-particles move as in a random barrier energy landscape. Moves of the zr particles cause changes of the center of mass coordinate as defined above: A zr-particle hopping to the right across a bond with $s_j > 0$ ($s_j < 0$) increases (decreases) the center of mass position by $1/L$. Conversely, a zr-particle hopping to the left across a bond with $s_j > 0$ ($s_j < 0$) decreases (increases) the center of mass position by $1/L$. As with the field free case of the open system, we use a quantum Hamiltonian and a tensor basis for the state space to compute the stationary state of the zr lattice of length M with a given $\mathbf{s}$ and $\mathbf{w}$. Let $e_1(i)$ denote an unoccupied site i and $e_m(i)$ where $m > 1$ a site i occupied by $n_i = m - 1$ particles. Here, vectors are infinite-dimensional. The matrix for creation of a particle at site i is then given by:

$$b_i^\dagger = \begin{pmatrix} 0 & 0 & \dots & 0 & \dots \\ 1 & 0 & \dots & 0 & \dots \\ \vdots & \ddots & \ddots & \vdots & \dots \\ 0 & & 1 & 0 & \dots \\ \vdots & \dots & \dots & \ddots & \ddots \end{pmatrix}_i. \quad (15)$$

The matrix for annihilation of a particle at site i is accordingly given by:

$$b_i = \begin{pmatrix} 0 & 1 & \dots & 0 & \dots \\ 0 & 0 & 1 & \dots & \dots \\ \vdots & \ddots & \ddots & \ddots & \dots \\ 0 & & 0 & 0 & \ddots \\ \vdots & \dots & \dots & \dots & \ddots \end{pmatrix}_i. \quad (16)$$

For constructing the diagonal part of the Hamiltonian we need the following type of

matrix (note that $\langle s|b_i^\dagger = \langle s|\mathbf{1}$ and $\langle s|b_i = \langle s|m_i$):

$$m_i = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots \\ 0 & 1 & \ddots & & \dots \\ \vdots & \ddots & \ddots & 0 & \dots \\ \vdots & & 0 & 1 & \ddots \\ \vdots & \dots & \dots & \ddots & \ddots \end{pmatrix}_i. \quad (17)$$

A hopping event from site i to site $i+1$ is described by the combined action of the matrices b_i and $b_{i+1}^\dagger$. This yields the following expression for the quantum Hamiltonian:

$$H_{zr} = \sum_{i=1}^M \left(-h_{i+1}b_{i+1}b_i^\dagger w_{i+1} - h_{i+1}^{-1}b_ib_{i+1}^\dagger w_{i+1} + h_{i+1}^{-1}w_{i+1}m_i + h_{i+1}w_{i+1}m_{i+1} \right). \quad (18)$$

A yet unnormalized product ansatz

$$|P_{zr}^* \rangle = \begin{pmatrix} 1 \\ z_1 \\ z_1^2 \\ \vdots \end{pmatrix} \otimes \dots \otimes \begin{pmatrix} 1 \\ z_M \\ z_M^2 \\ \vdots \end{pmatrix}. \quad (19)$$

yields

$$\begin{aligned} H_{zr}|P_{zr}^* \rangle &= \sum_{i=1}^M \left(-h_{i+1}w_{i+1}\frac{z_{i+1}}{z_i}m_i|P_{zr}^* \rangle - h_{i+1}^{-1}w_{i+1}\frac{z_i}{z_{i+1}}m_{i+1}|P_{zr}^* \rangle \right. \\ &\quad \left. + h_{i+1}^{-1}w_{i+1}m_i|P_{zr}^* \rangle + h_{i+1}w_{i+1}m_{i+1}|P_{zr}^* \rangle \right) \\ &= \sum_{i=1}^M \left(-h_{i+1}w_{i+1}\frac{z_{i+1}}{z_i}m_i|P_{zr}^* \rangle - h_i^{-1}w_i\frac{z_{i-1}}{z_i}m_i|P_{zr}^* \rangle \right. \\ &\quad \left. + h_{i+1}^{-1}w_{i+1}m_i|P_{zr}^* \rangle + h_iw_im_i|P_{zr}^* \rangle \right). \end{aligned} \quad (20)$$

$|P_{zr}^* \rangle$ is stationary provided

$$h_{i+1}w_{i+1}z_{i+1} + h_i^{-1}w_iz_{i-1} = h_{i+1}^{-1}w_{i+1}z_i + h_iw_iz_i. \quad (21)$$

Applying a general solution²⁴ to the case at hand and including normalization the stationary state for a channel characterized by $\mathbf{s}$ and $\mathbf{w}$ and with a total number of particles K yields

$$|P_{zr}^*(\mathbf{s}, \mathbf{w}, K, M) \rangle = \sum_{(n_1, \dots, n_M)} \prod_{i=1}^M z_i^{n_i} |n_1, \dots, n_M \rangle = \frac{1}{\sum_{(n_1, \dots, n_M)} \prod_{i=1}^M z_i^{n_i}}, \quad (22)$$

where

$$z_l = \sum_{i=1}^M \frac{1}{h_{l+i}^{-1} w_{l+i}} \prod_{j=1}^{i-1} h_{l+j}^2. \quad (23)$$

The primed sums are meant to be summations under the constraint $\sum_{i=1}^\sigma n_i = K$. Knowing the stationary state for each channel, the drift velocity for the individual channels can be computed. In the lattice gas picture, every time a particle $s_i = -1$ ($s_i = 1$) is hopping to the right (left) it changes the position of the center of mass by $1/(L+1)$. The opposite process, i.e. a particle $s_i = -1$ ($s_i = 1$) hopping to the left (right) changes the center of mass position by $-1/(L+1)$. Therefore, as mentioned above, $v = j^- - j^+$, i.e. the difference between the currents of negative and positive particles. This operator v translates to the zr picture as follows:

$$j^- - j^+ \rightarrow \frac{1}{L} \sum_{i=1}^M s_{i+1} w_{i+1} (h_{i+1}^{-1} b_i b_{i+1}^\dagger - h_{i+1} b_{i+1} b_i^\dagger). \quad (24)$$

Thus, the drift velocity of a given channel is given by

$$\begin{aligned} \langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle &= \langle s | \frac{1}{L} \sum_{i=1}^M (s_{i+1} h_{i+1}^{-1} w_{i+1} b_i b_{i+1}^\dagger \\ &\quad - s_{i+1} h_{i+1} w_{i+1} b_{i+1} b_i^\dagger) \Theta(K) | P_{zr}^* \rangle > \frac{1}{c_K} \end{aligned} \quad (25)$$

Here, $\Theta(K)$ projects on those states of $|P_{zr}^* \rangle$ which have a constant number of particles K and c_K is the normalization. So $\Theta(K)/c_K |P_{zr}^* \rangle = |P_{zr}^*(\mathbf{s}, \mathbf{w}, K) \rangle$. Using this form allows us to make use of the fact that due to the combined effect of $b_i b_j^\dagger$ which is to redistribute the particles without changing their number, we can commute $b_i b_j^\dagger$ and $\Theta(K)$. This allows us then to apply the matrices to the product measure, where its effect is easy to see.

$$\begin{aligned}
\langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle &= \langle s | \frac{1}{L} \sum_{i=1}^M (s_{i+1} h_{i+1}^{-1} w_{i+1} b_i b_{i+1}^\dagger \\
&\quad - s_{i+1} h_{i+1} w_{i+1} b_{i+1} b_i^\dagger) \Theta(K) | P_{zr}^* \rangle > \frac{1}{c_K} \\
&= \langle s | \frac{1}{L} \sum_{i=1}^M (s_{i+1} h_{i+1}^{-1} w_{i+1} \Theta(K) b_i b_{i+1}^\dagger \\
&\quad - s_{i+1} h_{i+1} w_{i+1} \Theta(K) b_{i+1} b_i^\dagger) | P_{zr}^* \rangle > \frac{1}{c_K} \\
&= \frac{1}{L} \sum_i (s_{i+1} h_{i+1}^{-1} w_{i+1} \frac{z_i}{z_{i+1}} \langle s | m_{i+1} | \Theta(K) P_{zr}^* \rangle \\
&\quad - s_{i+1} h_{i+1} w_{i+1} \frac{z_{i+1}}{z_i} \langle s | m_i | \Theta(K) P_{zr}^* \rangle) \frac{1}{c_K} \quad (26)
\end{aligned}$$

We now have to calculate the expression c_K for the product state:

$$c_K = \sum_{(n_1, \dots, n_M)}', \prod_{i=1}^M z_i^{n_i} \quad (27)$$

where the primed sum again means summing with the constraint $\sum_{i=1}^M n_i = K$.

A similar type of summation is found when regarding the sums of the type

$$\langle s | m_\beta \frac{\Theta(K)}{c_K} | P_{zr}^* \rangle = \sum_{(n_1, \dots, n_M)}'' \frac{1}{c_K} \prod_{j=1}^M z_j^{n_j}, \quad (28)$$

where the double primed sum has the constraints $\sum_{j=1}^M n_j = K$ and $n_\beta \neq 0$. A straightforward calculation of the sums is impossible due to the constraints. To simplify the task, we can profitably use the following identity¹²:

$$\sum_{(n_1, \dots, n_M)}', \prod_{i=1}^M z_i^{n_i} = \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^{K+1}} \sum_{(n_1, \dots, n_M)} \prod_{j=1}^M (\alpha z_j)^{n_j}. \quad (29)$$

Thus we can transform the sums into integrals:

$$\begin{aligned}
c_K &= \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^{K+1}} \sum_{(n_1, \dots, n_M)} \prod_{j=1}^M (\alpha z_j)^{n_j} \\
&= \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^{K+1}} \prod_{j=1}^M \frac{1}{1 - \alpha z_j} \\
&= Q_{K,M}, \quad (30)
\end{aligned}$$

$$\begin{aligned}
\langle s | m_\beta \frac{\Theta(K)}{c_K} | P_{zr}^* \rangle &= \frac{1}{c_K} \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^{K+1}} \sum_{(n_1, \dots, n_M), n_\beta \neq 0} \prod_{j=1}^M (\alpha z_j)^{n_j} \\
&= \frac{1}{c_K} z_\beta \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^K} \prod_{j=1}^M \frac{1}{1 - \alpha z_j} \\
&= z_\beta \frac{Q_{K-1, M}}{Q_{K, M}}.
\end{aligned} \tag{31}$$

The integrals $Q_{K, M}$ satisfy the recursion relation

$$Q_{K, M} = Q_{K, M-1} + z_M Q_{K-1, M}. \tag{32}$$

An explicit solution of this relation yields:

$$Q_{K, M} = \sum_{i=1}^M z_i^{K+M-1} \prod_{l=1}^M (z_i - z_l)^{-1}. \tag{33}$$

The primed product indicates $i \neq l$.

From this point on we are not going to carry through the complete calculation for $\langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle$. We will expand it into a series in E and keep only first order terms which is sufficient for employing the Einstein relation. We start from the expression for $\langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle$ after having inserted the result for $\langle s | m_i \frac{\Theta(N-L)}{c_K} | P_{zr}^* \rangle$:

$$\begin{aligned}
\langle v(s, w, K) \rangle &= \frac{1}{L} \sum_{i=1}^M (s_{i+1} h_{i+1}^{-1} w_{i+1} \frac{z_i}{z_{i+1}} z_{i+1} \frac{Q_{K-1, M}}{Q_{K, M}} \\
&\quad - s_{i+1} h_{i+1} w_{i+1} \frac{z_{i+1}}{z_i} z_i \frac{Q_{K-1, M}}{Q_{K, M}}) \\
&= \frac{1}{L} \sum_{i=1}^M s_{i+1} (h_{i+1}^{-1} w_{i+1} z_i - h_{i+1} w_{i+1} z_{i+1}) \frac{Q_{K-1, M}}{Q_{K, M}} \\
&= \frac{1}{L} \sum_{i=1}^M s_{i+1} (1 - \exp(-ES)) \frac{Q_{K-1, M}}{Q_{K, M}} \\
&= \frac{S}{L} (1 - \exp(-ES)) \frac{Q_{K-1, M}}{Q_{K, M}},
\end{aligned} \tag{34}$$

where $S = \sum_{i=1}^M s_i$. For the second equality we used the explicit form of the z_i . The term in brackets is easily expanded into a series in E : $1 - \exp(ES) = ES + o(E^2)$. We keep only the first order term in E . This means that when we are expanding the expression

$Q_{K-1,M}/Q_{K,M}$ we need only to keep zero order terms in E , as all other contributions will vanish when using the Einstein relation. The only terms containing E in the $Q_{K,M}$ are the z_i , for which we find to first order in E :

$$\begin{aligned}
z_i &= \sum_{j=1}^M \frac{1}{h_{i+j}^{-1} w_{i+j}} \prod_{k=1}^{j-1} h_{i+k}^2 \\
&= \sum_{j=1}^M \frac{1}{w_{i+j}} \left(1 - \frac{E}{2} s_{j+i} + \dots\right) \prod_{k=1}^{j-1} (1 - E s_{i+k} + \dots) \\
&= \sum_{j=1}^M \frac{1}{w_{i+j}} + o(E). \\
&= z + o(E).
\end{aligned} \tag{35}$$

Here, z is defined as the zero field value of z_i . The fact that all z_i are equal at zero field is decisive for the explicit integration of $Q_{K,M}$. We obtain:

$$\begin{aligned}
Q_{K,M} &= \frac{1}{2\pi i} \oint \frac{d\alpha}{\alpha^{K+1}} \prod_{j=1}^M \frac{1}{1 - \alpha z_j} \\
&= \binom{M+K-1}{M-1} z^K.
\end{aligned} \tag{36}$$

It is now clear how to proceed with evaluating $\langle v(s, w, K, M) \rangle$, as

$$\frac{Q_{K-1,M}}{Q_{K,M}} = \frac{\binom{K+M-2}{M-1} z^{M-1}}{\binom{K+M-1}{M-1} z^K} = \frac{K}{K+M-1} \frac{1}{z}. \tag{37}$$

This yields for $\langle v(s, w, K, M) \rangle$:

$$\begin{aligned}
\langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle &= \frac{1}{L} \sum_{i=1}^M s_{i+1} E S \frac{K}{K+M-1} \frac{1}{z} + o(E^2) \\
&= \frac{1}{L} E S^2 \frac{K}{K+M-1} \frac{1}{z} + o(E^2).
\end{aligned} \tag{38}$$

This is our final expression for $\langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle$. Knowing the drift velocity for each channel characterized by $\mathbf{s}, \mathbf{w}$ we have to give each of those a weight $\Psi(\mathbf{s}, \mathbf{w}, K, M)$ when averaging over the channels. Following an argument by Prähofer²², we show below that provided a configuration in the zr picture is weighted such that at $E = 0$ its probability as contained in $P_{ring}^*(\mathbf{s}, \mathbf{w}, K, M) = \Psi(\mathbf{s}, \mathbf{w}, K, M) P_{zr}^*(\mathbf{s}, \mathbf{w}, K, M)$ is equal to the probability

of the corresponding state in the lattice gas of the open system as given by $P_{open}^*(0)$, for the diffusion constants the following relation holds: $D_{L+1}^{open} \geq D_L^{per}$.

When relating zr to lattice gas probabilities it has to be kept in mind that the periodic zr system has M sites, while the corresponding periodic lattice gas has $L \geq M$ sites. This means a state of the zr system with probability q_{zr} can be permuted M times where due to translational invariance all resulting states have the same probability. A lattice gas configuration with probability q_{lg} can be cycled through L permutations leading to equal probabilities. Therefore, equal probabilities of respective configurations means: $Mq_{zr} = Lq_{lg}$. The weight factor thus has to be chosen as follows:

$$\Psi(\mathbf{s}, \mathbf{w}, K, M) = \frac{d^M}{(2d+1)^L} \binom{L}{M} \prod_{j=1}^M f(w_j). \quad (39)$$

Thus we find for the drift velocity when averaged over the channels:

$$\begin{aligned} \bar{v} &= \sum_M \sum_{\mathbf{s}=(s_1, \dots, s_M)} \sum_{\mathbf{w}=(w_1, \dots, w_M)} \Psi(\mathbf{s}, \mathbf{w}) \langle v(\mathbf{s}, \mathbf{w}, K, M) \rangle \\ &= \sum_M \frac{1}{L} EM \frac{d^M}{(2d+1)^L} \binom{L}{M} \frac{L-M}{L-1} \left(\sum_w \prod_{j=1}^M f(w_j) \frac{1}{z} \right). \end{aligned} \quad (40)$$

This can be rewritten as

$$\bar{v} = E \sum_{M=1}^L \Omega(M) M \left\langle \frac{1}{z} \right\rangle. \quad (41)$$

Here $\Omega(M)$ contains all the factors depending on M occurring in the previous expression and $\langle 1/z \rangle$ is a disorder average. $\Omega(M)$ is a function that is sharply peaked at $M = L/2$ implying that in the limit for large L only terms with large M significantly contribute to the result. In the limiting case of $L \rightarrow \infty$ invoking the central limit theorem (remember that we demanded $\langle 1/W \rangle$ and $\langle 1/W^2 \rangle$ to be finite) yields:

$$\left\langle \frac{1}{z} \right\rangle = \frac{1}{M \langle 1/W \rangle}. \quad (42)$$

Therefore

$$\bar{v} = \sum_{M=1}^L \frac{1}{L} EM \frac{d^M}{(2d+1)^L} \binom{L}{M} \frac{L-M}{L-1} \frac{1}{M} \frac{1}{\langle \frac{1}{W} \rangle}. \quad (43)$$

Using the Einstein relation yields in the limit for large L

$$\begin{aligned} D(0) &= \frac{1}{L(L-1)} \frac{1}{(2d+1)} \left(1 - \frac{1}{(2d+1)^{L-1}}\right) \frac{1}{\langle \frac{1}{W} \rangle} \\ &= D^* \frac{1}{\langle \frac{1}{W} \rangle}, \end{aligned} \tag{44}$$

where D^* is the diffusion constant for the ordered case^{12,13}.

Thus we have shown that $\lim_{L \rightarrow \infty} DL^2 = 1/[(2d+1) \langle 1/W \rangle]$ for the periodic case, which is a lower bound to the open case. Note that the factor $1/\langle 1/W \rangle$ is the same as occurring in the single particle diffusion constant for the random barrier model. In this model random energy barriers are assigned to bonds between sites just as in the zr picture of our problem.

IV. MODEL WITH OPEN BOUNDARY CONDITIONS

A. Variational formula

In order to find an upper bound on the diffusion constant we follow the strategy of Ref. 13. The quantum Hamiltonian H_{open} for our model in its representation using the tensor basis can be decomposed into the sum of a diagonal part (D) and a non diagonal one (M). Each of these can themselves be split into a part describing those moves leading to an increase of the center of mass coordinate (M^+ and D^+) and a respective part connected to a decrease (M^- and D^-). Thus $H = D^+ + D^- - M^+ - M^-$.

The following statement²¹ holds for the diffusion constant D :

$$\begin{aligned} D &= \inf_{g \in \Omega} \left[\frac{1}{2(L+1)^2} \langle D^+ + D^- \rangle - \frac{2}{(L+1)} \langle s|[D^+ - D^-]g|P^* \rangle + \langle s|gHg|P^* \rangle \right] \\ &= \inf_{g \in \Omega} F[g]. \end{aligned} \tag{45}$$

Here, Ω is the space of diagonal matrices with dimension $(2\sigma+1)^L$. Thus plugging in any diagonal matrix g into the functional $F[g]$ yields an upper bound on D . The challenge is to choose g such that the upper bound gets as small as possible. Due to the dimensionality of the diagonal matrices increasing exponentially with L , an exact minimization is not feasible.

Still, some information can be gained by observing that the functional is convex and thus the minimizing $g_0 \in \Omega$ is the only matrix for which $\delta F[g]$ vanishes. Choosing g_0 such that

$$\langle s | g_0 H = \langle s | [D^+ - D^-] \quad (46)$$

the variation vanishes. Unfortunately, this formula cannot be solved for g_0 . We will show below how still some insight can be gained from this equation. Introducing the matrix d , where $d(y'|y) = \langle y' | d | y \rangle$ gives the change in the CMS coordinate when a transition from state y to y' is made and the matrix w , where $w(y'|y)$ denotes the corresponding rate, the variational formula may be written as:

$$D = \inf \frac{1}{2} \sum_{y', y \in X} w(y'|y) P^*(y) [d(y'|y) + g(y') - g(y)]^2, \quad (47)$$

where $g(y) = \langle s | g | y \rangle$. Following Ref. 22 we now prove that $D_L^{open} \geq D_{L+1}^{per}$. To simplify the notation we remark that in our model with periodic boundary conditions as in the original RD model transitions from a given state y are only allowed to a state $y' = y^{i,i+1}$, where the spins y_i and y_{i+1} are interchanged. Thus the last formula applied to the periodic case reads as

$$D_{L+1}^{per} = \inf \frac{1}{2} \sum_{l=0}^L \left(\sum_{y \in X} w(y^{l,l+1} | y) P^*(y) [d(y^{l,l+1} | y) + g(y^{l,l+1}) - g(y)]^2 \right). \quad (48)$$

Here, each transition between states in a lattice gas of length $L + 1$ changes the cm coordinate by $\pm 1/(L + 1)$, as there are as many reptons as bonds between them. In contrast to that in the open boundary case there is one repton more as there are bonds leading to the cm coordinate changing by $\pm 1/(L + 1)$ in any transition of a lattice gas of length L . Here, transitions between states can not only occur by interchanging spins y_1 and y_{i+1} in the bulk, but also by creation and annihilation events at the ends. These events can equivalently be viewed as interchanges of the spins y_1 or y_L respectively with 'imaginary' spins y_0 and y_{L+1} provided the rates for these interchanges at the ends are adapted such that they correspond to the creation and annihilation rates as demanded by the model. Taking the distribution at the 'imaginary' sites as for any other site in the lattice gas (remember that we have a

homogeneous product measure) the introduction of a parameter

$$c_l = \begin{cases} 2d + 1 & \text{if } l = 0 \text{ or } l = L \\ 1 & \text{for } l = 1, \dots, L - 1 \end{cases} \quad (49)$$

leads to a fulfillment of this requirement and the functional for the diffusion constant of the open chain of length L reads as

$$D_L^{open} = \inf \frac{1}{2} \sum_{l=0}^L c_l \left(\sum_{y \in X} w(y^{l,l+1}|y) P^*(y) [d(y^{l,l+1}|y) + g(y^{l,l+1}) - g(y)]^2 \right) \quad (50)$$

where w and d are as for the periodic case. Letting $\mathbf{y} = (y_0, \dots, y_{L+1})$ for open and periodic system and observing that as y_0 does not occur in the functional for D_{L+1}^{per} the averaging of y_0 over P^* yields 1 and thus does not change the result.

$$\begin{aligned} & D_L^{open}[g] - D_{L+1}^{per}[g] \\ &= c_0 \left(\sum_{y \in X} w(y^{0,1}|y) P^*(y) [d(y^{0,1}|y) + g(y^{0,1}) - g(y)]^2 \right) \\ & \quad + c_L \left(\sum_{y \in X} w(y^{L,L+1}|y) P^*(y) [d(y^{L,L+1}|y) + g(y^{L,L+1}) - g(y)]^2 \right) \\ & \quad - \sum_{y \in X} w(y^{L+1,1}|y) P^*(y) [d(y^{0,1}|y) + g(y^{0,1}) - g(y)]^2 \\ & \quad - \sum_{y \in X} w(y^{L,L+1}|y) P^*(y) [d(y^{L,L+1}|y) + g(y^{L,L+1}) - g(y)]^2 \\ &= 2 \sum_{y \in X} w(y^{L+1,1}|y) P^*(y) [d(y^{0,1}|y) + g(y^{0,1}) - g(y)]^2 \\ & \quad + 2 \sum_{y \in X} w(y^{L,L+1}|y) P^*(y) [d(y^{L,L+1}|y) + g(y^{L,L+1}) - g(y)]^2 \\ & \geq 0 \end{aligned} \quad (51)$$

Therefore, the diffusion constant of the periodic chain with length $L + 1$ is a lower bound for the open chain with length L .

B. Functional for diffusion with kinematic disorder

The space of diagonal matrices Ω from which an appropriate $g \in \Omega$ has to be chosen such that $F[g]$ gets as small as possible, has dimension $(2\sigma + 1)^L$. A scalar product on Ω

can be defined for arbitrary $m, n \in \Omega$ as $\langle s|mn|P^* \rangle$. A basis for Ω can be built from the matrices $\hat{y}_1(i)$ to $\hat{y}_{2\sigma}(i)$ acting non trivially only at site i and the unit matrix. We specify only the following:

$$\hat{y}_\alpha(i) = \frac{1}{2d+1}E_{(2\alpha,2\alpha)}(i) - \frac{1}{2d+1}E_{(2\alpha+1,2\alpha+1)}(i) \text{ for } \alpha = 1, \dots, \sigma, \quad (52)$$

$$\hat{y}_{\sigma+1}(i) = -\frac{2d}{2d+1}E_{(1,1)}(i) + \sum_{j=2}^{2\sigma+1} \frac{1}{2d+1}E_{(j,j)}(i). \quad (53)$$

For the remaining $\sigma - 1$ matrices we demand that they are chosen such that for all $a, b = 1, \dots, 2\sigma$, $i, j = 1, \dots, L$ and $a \neq b$ we have $\langle \hat{y}_a(i)\hat{y}_b(j) \rangle = 0$ as well as $\langle \hat{y}_a(i) \rangle = 0$. It is easily checked that the defined matrices are mutually orthogonal. A basis for the space of diagonal matrices Ω is then given by:

$$\left\{ \prod_{j=1}^{2\sigma+1} \hat{y}_j(I_j) | I_j \subset \{1, \dots, L\} \quad \forall j; I_\alpha \cap I_\beta = \emptyset \quad \forall \alpha \neq \beta \right\}. \quad (54)$$

The choice of a $g \in \Omega$ fulfilling $\langle s|[D^+ - D^-] = \langle s|gH$ leads to $\delta F[g] = 0$. In spite of this relation not being solvable for g information can be gained by observing that Ω contains subsets $\Omega_{i_1, \dots, i_\sigma}$ which are invariant to H in the sense that if $g \in \Omega_{i_1, \dots, i_\sigma}$, then the diagonal matrix g' fulfilling $\langle s|gH = \langle s|g'$ is in $\Omega_{i_1, \dots, i_\sigma}$ too, where

$$\Omega_{i_1, \dots, i_\sigma} = \text{span} \left\{ \prod_{j=1}^{2\sigma+1} \hat{y}_j(I_j) | I_j \subset \{1, \dots, L\} \quad \forall j; I_\alpha \cap I_\beta = \emptyset \quad \forall \alpha \neq \beta; |I_j| = i_j \quad \forall j = 1, \dots, \sigma \right\}. \quad (55)$$

By calculating $\langle s|[D^+ - D^-]$ we find that the minimizing g_0 is in the subspace $\bigoplus_{j=1}^{\sigma} \Omega_{i_1, \dots, i_\sigma}$ where $i_k = \delta_{i,k} \quad \forall \quad k = 1, \dots, \sigma$. Thus the most general ansatz for g based on this information is:

$$g = \sum_{\alpha=1}^{\sigma} \sum_{i=0}^{L+1} \sum_{I_{\sigma+1}, \dots, I_{2\sigma+1}} c_{i, I_{\sigma+1}, \dots, I_{2\sigma+1}}^{\alpha} \hat{y}_\alpha(i) \hat{y}_{\sigma+1}(I_{\sigma+1}) \dots \hat{y}_{2\sigma+1}(I_{2\sigma+1}) \quad (56)$$

where the coefficients c are real numbers and the sets $I_\sigma, \dots, I_{2\sigma+1}$ are mutually disjoint.

C. Ansatz for g

The general ansatz given above for g contains too many parameters for letting a minimization appear feasible. Instead we use a generalization of the ansatz used by Prähofer²² for the ordered RD model:

$$g = \sum_{\alpha=1}^{\sigma} \left\{ \sum_{i=0}^{L+1} a_i^{\alpha} \hat{y}_{\alpha}(i) + \sum_{k,k'=0}^{L+1} c_{k,k'}^{\alpha} \hat{y}_{\alpha}(k) \hat{y}_{\sigma+1}(k') \right. \\ \left. + \sum_{o,p,q=0;p < q}^{L+1} e_{o,p,q}^{\alpha} \hat{y}_{\alpha}(o) \hat{y}_{\sigma+1}(p) \hat{y}_{\sigma+1}(q) \right\}. \quad (57)$$

The primed sums indicate the summation variables to be mutually different. Note that this ansatz reflects a particular choice of parameters for the general ansatz. Inserting the trial function into $F[g]$ yields the following functional:

$$D[g] \\ = \sum_{\alpha=1}^{\sigma} \left\{ \frac{2d}{(2d+1)^2} f(W_{\alpha}) W_{\alpha} \sum_{l=0}^L c_l \right. \\ \left[\left(\frac{1}{L+1} + (a_l^{\alpha} - a_{l+1}^{\alpha}) - \frac{2d}{(2d+1)} (c_{l,l+1}^{\alpha} - c_{l+1,l}^{\alpha}) \right)^2 \right. \\ \left. + \frac{2d}{(2d+1)^2} \sum_{k'} \left(c_{l,k'}^{\alpha} - c_{l+1,k'}^{\alpha} - \frac{2d}{(2d+1)} (e_{l,(l+1,k')}^{\alpha} - e_{l+1,(l,k')}^{\alpha}) \right)^2 \right. \\ \left. + \sum_{\beta=1}^{\sigma} \frac{2d}{(2d+1)} f(W_{\beta}) \sum_k \left(c_{k,l}^{\beta} - c_{k,l+1}^{\beta} \right)^2 \right. \\ \left. + \frac{4d^2}{(2d+1)^4} \sum_{p < q} \left(e_{l,p,q}^{\alpha} - e_{l+1,p,q}^{\alpha} \right)^2 \right. \\ \left. + \sum_{\beta=1}^{\sigma} \frac{4d^2}{(2d+1)^3} f(W_{\beta}) \sum_{o,q} \left(e_{o,(l,q)}^{\beta} - e_{o,(l+1,q)}^{\beta} \right)^2 \right] \left. \right\}. \quad (58)$$

The free parameters are the coefficients a_i^{α} , $c_{k,k'}^{\alpha}$ and $e_{o,p,q}^{\alpha}$ which are, generalizing results in Ref. 13 chosen as

$$a_{i+1}^{\alpha} = \sum_{l=0}^i \left(\frac{1}{L+1} - \frac{2d}{(2d+1)} (c_{l,l+1}^{\alpha} - c_{l+1,l}^{\alpha}) \right) \\ a_0^{\alpha} = 0 \quad (59)$$

$$c_{i,j}^\alpha = \begin{cases} \frac{(2d+1)C}{2d} (1 - g(\frac{i}{n})) \frac{L+1-j}{(L+1)^2} & \text{for } i < j \\ -c_{L+1-i, L+1-j}^\alpha & \text{for } i > j \end{cases} \quad (60)$$

$$e_{o,p,q}^\alpha = \begin{cases} \frac{(2d+1)^2 C}{4dn} g'(\frac{o}{n}) \frac{(L+1-p)(L+1-q)}{(L+1)^2 (L-o)} & \text{for } o < p < q \\ -e_{L+1-o, L+1-q, L+1-p}^\alpha & \text{for } p < q < o. \end{cases} \quad (61)$$

Here, g is a monotonically decreasing smooth function on the real numbers with $g(x) = 1$ for $x \leq 0$ which is decreasing exponentially fast for $x \rightarrow \infty$. This function has the property that

$$\sum_{i=0}^{\infty} \left[g^{(r)} \left(\frac{i}{n} \right) \right]^s = O(n), \quad (62)$$

where r, s are integers, and $n = (L+1)^{0.75}$. Given this choice, the following upper bound for D_{L+1}^{open} is found

$$D_{L+1}^{open} \leq \frac{1}{(2d+1)} \frac{1}{(L+1)^2} < W >, \quad (63)$$

where $< W >$ is the disorder average.

This completes the proof that asymptotically $D \propto 1/L^2$ in the presence of kinematic disorder.

V. MC RESULTS

In the case of the ordered RD model the results for D in the periodic case¹² coincide to leading order with the upper bound for the open case¹³. In our model we managed to give bounds on D which both scale with $1/L^2$ in the limit for long chains. This implies that kinematic disorder does not ruin the scaling relation of standard reptation theory. Still, the numerical prefactors differ. We believe that it is the result for the lower bound which correctly describes the diffusive behavior for long chains. This lower bound was obtained by a rigorous calculation while the upper bound results from a variational treatment. Given that our conjecture is true and for long chains $D = 1/[L^2(2d+1) < 1/W >]$ then for any choice of disorder distribution $DL^2 < 1/W >$ plotted against the chain length is constant. We performed MC simulations with different disorder distributions. It turns out that $L = 30$

is sufficient to observe the curves all converging to the same constant value. For the cases $\langle 1/W \rangle = 1$, $\langle 1/W \rangle = 0.02$ and $\langle 1/W \rangle = 0.03$ figure 3 shows our MC results. These results suggest that in the long chain length limit $D = 1/[L^2(2d + 1) \langle 1/W \rangle]$.




Figure 3: $D(0)L^2 \langle 1/W \rangle$ versus L for the cases as described in the text.

VI. DYNAMICS OF INTERNAL SEGMENTS

The surprising result of the previous sections is the observation that the effect of kinematic disorder on the collective behavior of all connected polymer segments is (to leading order in system size) the same as on a simple point like object in the same disorder environment. In order to understand this observation we now consider the dynamics of the internal segments in the hydrodynamic limit of vanishing lattice spacing. For the local concentration of reptons one obtains from the usual ordered repton model in this limit Rouse dynamics¹⁵ restricted to motion inside the tube⁵. The boundary dynamics, i.e., the hopping into and out of the tube resp. describe the entropic tensile force acting on the chain ends and keeping the polymer in its stretched equilibrium conformation. For understanding the dependence of the diffusion coefficient on kinematic disorder, which affects mostly the bulk of the polymer chain, it is sufficient to focus on the hydrodynamic behavior of the bulk reptons. In order

to study this limit it is convenient to first investigate the associated zero-range process and then translate the result into the hydrodynamic limit of the exclusion process.

For the zero-range process described in Section III with $E = 0$, the average number of particles $\rho_i(t)$ at site i obeys the exact time evolution equation given by

$$\frac{\partial \rho_i(t)}{\partial t} = W_i z_{i-1}(t) + W_{i+1} z_{i+1}(t) - (W_i + W_{i+1}) z_i(t) \quad (64)$$

where $z_i(t)$ is the probability that site i is occupied at time t . It is known that in the steady state, the occupancy probability z_i^{ss} is spatially uniform and can be related to the steady state particle density as¹⁸

$$z_i^{ss} = \frac{\rho_i^{ss}}{1 + \rho_i^{ss}}. \quad (65)$$

It follows that the steady state density profile is uniform in spite of disorder.

To understand the dynamics in the hydrodynamic limit of vanishing lattice spacing, we expand Eq.(64) to second order in lattice constant and find

$$\frac{\partial \rho(x, t)}{\partial t} = \frac{\partial}{\partial x} \left[W(x) \frac{\partial z(x, t)}{\partial x} \right] \quad (66)$$

At large enough time, the system is expected to be in local equilibrium so that we may assume the usual approximation for $z(x, t) - \rho(x, t)$ in the steady state to be valid: $z(x, t) \approx \rho(x, t)/(1 + \rho(x, t))$, where equation (65) was used. We are interested in the density fluctuations about the steady state, $\Delta \rho(x, t) = \rho(x, t) - \rho^{ss}(x)$ where $\rho^{ss}(x) = \rho = K/M$. Retaining the lowest nonvanishing term in the expansion of the preceding equation in powers of $\Delta \rho(x, t)$, we obtain

$$\frac{\partial \Delta \rho(x, t)}{\partial t} = \frac{1}{(1 + \rho)^2} \frac{\partial}{\partial x} \left[W(x) \frac{\partial \Delta \rho(x, t)}{\partial x} \right] \quad (67)$$

The above equation describes a random walker in one dimension, diffusing in a random medium with bond-symmetric hopping rates $W(x)$. It can be shown that at large time and length scales, the random walker can be described by a single, effective diffusion constant, $\mathcal{D} = 1/[(1 + \rho)^2 \langle 1/W \rangle]$, provided $\langle 1/W \rangle$ is finite²⁵. Thus we obtain the bulk diffusion constant to be given by $\mathcal{D}$ in the zero range process.

Regarding the site as particle and mass as hole clusters, the above model maps onto symmetric exclusion process (SEP) with particlewise disorder. We want to calculate the bulk diffusion constant in the SEP picture using the above results for zero range process. Since the steady state density profile is uniform in both the pictures, the average local

density n_i in the vicinity of the location of particle i in SEP is related to that in the zero range process as $n_i = 1/(1 + \rho_i)$. Then the density fluctuation about the steady state in the vicinity of particle i is given by

$$\Delta n_i = \frac{-1}{(1 + \rho)^2} \Delta \rho_i + O((\Delta \rho)^2) \quad . \quad (68)$$

We note that the above is true only at large enough times, as was pointed out in a similar analysis for the tagged particle correlation function for SEP without disorder²⁶.

Using Eq.(67) and Eq.(68), we find that

$$\frac{\partial \Delta n(x, t)}{\partial t} = \frac{1}{(1 + \rho)^2} \frac{\partial}{\partial x} \left[W(x) \frac{\partial \Delta n(x, t)}{\partial x} \right] \quad . \quad (69)$$

Further note that x is the space index in the zero range process, while it labels the particles in the SEP. The space coordinate y in symmetric exclusion process is related to x as

$$y \approx \int^x dx' \rho(x') + x \quad , \quad (70)$$

which gives

$$\frac{\partial}{\partial x} = (1 + \rho) \frac{\partial}{\partial y} + O(\Delta n) \quad . \quad (71)$$

Thus the particle density fluctuations in the SEP obey

$$\frac{\partial \Delta n(y, t)}{\partial t} = \frac{\partial}{\partial y} \left[W(y) \frac{\partial \Delta n(y, t)}{\partial y} \right] \quad . \quad (72)$$

Using the random walker analogy, we obtain the effective diffusion constant $\overline{D}$ at large times to be equal to $1/\langle 1/W \rangle$ in the symmetric exclusion process.

This calculation shows that the internal segments of the polymer chain perform Rouse-type dynamics also in the presence of kinematic disorder, but with a disorder-dependent diffusion coefficient. This explains the occurrence of the same correction to the diffusion coefficient for the long time behavior of the polymer chain as a whole.

VII. CONCLUSIONS

It is the aim of this paper to disentangle the effects of the various types of disorder which one may expect to have significant impact on the dynamics of systems of entangled flexible polymers. We have focused on kinematic disorder which leaves the equilibrium conformation unchanged compared to a hypothetical ordered entanglement network (which

could in principle be manufactured artificially by placing a single polymer in an ordered array of obstacles on a surface). We have proved (rigorously in terms of the RD-model for reptation) that the asymptotic length scaling of the diffusion coefficient of the polymer chain remains as predicted by standard reptation theory. By studying the hydrodynamic limit we have shown that the individual polymer segments inside the tube perform Rouse dynamics in a disordered environment which corresponds to a system of local random barriers. Therefore the amplitude of the diffusion coefficient becomes dependent on the disorder in the same way a single particle in a random barrier system.

VIII. ACKNOWLEDGMENT

K.J. wishes to thank the Forschungszentrum Jülich for kind hospitality. R.D.W. thanks Deutsche Forschungsgemeinschaft for financial support.

* Electronic address: r.willmann@fz-juelich.de

- ¹ P. G. de Gennes, *Scaling concepts in polymer physics* (Cornell University Press, Ithaca, 1979).
- ² T. P. Lodge, Phys. Rev. Lett. **83**, 3218 (1999).
- ³ T. P. Lodge and E. D. von Meerwall, Macromolecules **33**, 1747 (2000).
- ⁴ J. D. Ferry, *Viscoelastic Properties of Polymers* (Wiley, New York, 1980).
- ⁵ M. Doi, *Introduction to Polymer Physics* (Oxford University Press, 1997).
- ⁶ E. Carlon, A. Drzewinski and J. M. J. van Leeuwen, Phys. Rev. E **64**, 010801(R) (2001).
- ⁷ M. Paessens and G. M. Schütz, Phys. Rev. E **66**, 021806 (2002).
- ⁸ M. Rubinstein, Phys. Rev. Lett. **59**, 1946 (1987).
- ⁹ J. M. Deutsch and T. L. Madden, J. Chem. Phys. **91**, 3252 (1989).
- ¹⁰ T. A. J. Duke, Phys. Rev. Lett. **62**, 2877 (1989).
- ¹¹ T. A. J. Duke, J. Chem. Phys. **93**, 9049 (1990).
- ¹² J. M. J. van Leeuwen and A. Kooiman, Physica A **184**, 79 (1992).
- ¹³ M. Prähofer and H. Spohn, Physica A **233**, 191 (1996).
- ¹⁴ G. T. Barkema, C. Caron, and J. F. Marko, Biopolymers **38**, 665 (1996).
- ¹⁵ G. M. Schütz, Europhys. Lett **48**, 623 (1999).

- ¹⁶ A. Baumgärtner and M. Muthukumar in *Advances in Chemical Physics, Volume XCIV*, edited by I. Prigogine and S. A. Rice (Wiley, New York, 1996).
- ¹⁷ A. E. Likhtman, S. T. Milner, and T. C. B. McLeish, Phys. Rev. Lett. **85**, 4550 (2000).
- ¹⁸ R. D. Willmann, J. Chem. Phys. **116**, 2688 (2002).
- ¹⁹ L. Schäfer, A. Baumgärtner, and U. Ebert, Eur. Phys. J. B **10**, 105 (1999).
- ²⁰ G. M. Schütz, in *Phase Transitions and Critical Phenomena*, Vol.19, edited by C. Domb and J. L. Lebowitz (Academic, London, 2000).
- ²¹ H. Spohn, *Large Scale Dynamics of Interacting Particles* (Springer, Heidelberg, 1991).
- ²² M. Prähofer, Diplom thesis, LMU Munich, 1994.
- ²³ I. Benjamini, P. A. Ferrari, and C. Landim, Stoch. Proc. Appl. **61**, 181 (1996).
- ²⁴ B. Derrida, J. Stat. Phys. **31**, 433 (1983).
- ²⁵ K. W. Kehr, K. Mussawisade, G. M. Schütz and T. Wichmann in *Diffusion in Condensed Matter* edited by P. Heitjans and J. Kräger (Springer, Berlin, 2003).
- ²⁶ S. Alexander and P. Pincus, Phys. Rev. B **18**, 2011 (1978).